

Mild traumatic brain injury recovery: a growth curve modelling analysis over two years

Ellen L. Carroll,¹ Joanne G. Outtrim,¹ Faye Forsyth,¹ Anne E. Manktelow,¹ Peter J. A. Hutchinson,² Olli Tenovuo,^{3,4} Jussi P. Posti,^{3,4,5} Lindsay Wilson,⁶ Barbara J. Sahakian,^{7,8,9} David K. Menon,^{1,9} & Virginia F. J. Newcombe.¹

- ¹ Division of Anaesthesia, Department of Medicine, University of Cambridge, UK
- ² Division of Neurosurgery, Department of Clinical Neurosciences, University of Cambridge, UK
- ³ Turku Brain Injury Centre, Turku University Hospital, Turku, Finland
- ⁴ Department of Clinical Neurosciences, University of Turku, Turku, Finland
- ⁵ Neurocenter, Department of Neurosurgery, Turku University Hospital, Turku, Finland
- ⁶ Division of Psychology, University of Stirling, Stirling, UK
- ⁷ Department of Psychiatry, University of Cambridge, UK
- ⁸ Behavioural & Clinical Neuroscience Institute, University of Cambridge, UK
- ⁹ Wolfson Brain Imaging Centre, Department of Clinical Neurosciences, University of Cambridge, UK

Corresponding author: Virginia F. J. Newcombe; Address: University Division of Anaesthesia, Box 93, Addenbrooke's Hospital, Hills Road, Cambridge, CB2 0QQ, UK; Email: vfjn2@cam.ac.uk; Telephone: +44 1223 586742; Fax: +44 1223 217887
Orcid id: <https://orcid.org/0000-0001-6044-9035>

Number of characters in the title: 91

Number of characters in the running head: 49

Key words: Mild traumatic brain injury; latent growth curve modelling; cognitive impairment; outcome.

ABSTRACT

Background: An improved understanding of the trajectory of recovery after mild traumatic brain injury is important to be able to understand individual patient outcomes, for longitudinal patient care and to aid the design of clinical trials.

Objective: To explore changes in health, well-being and cognition over the two years following mTBI using latent growth curve (LGC) modelling.

Methods: Sixty-one adults with mTBI presenting to a UK Major Trauma Centre completed comprehensive longitudinal assessment at up to five time points after injury; two weeks, three months, six months, one year and two years.

Results: Persisting problems were seen with neurological symptoms, cognitive issues and poor quality of life measures including 28% reporting incomplete recovery on the Glasgow Outcome Score Extended at two years. Harmful drinking, depression, psychological distress, disability, episodic memory, and working memory did not improve significantly over the two years following injury. For other measures, including the Rivermead Post-Concussion Symptoms and Quality of Life after Brain Injury (QOLIBRI), LGC analysis revealed significant improvement over time with recovery tending to plateau at three to six months.

Interpretation: Significant impairment may persist as late as two years after mTBI despite some recovery over time. Longitudinal analyses which make use of all available data indicate that recovery from mTBI occurs over a longer timescale than is commonly believed. These findings point to the need for long-term management of mTBI targeting individuals with persisting impairment.

INTRODUCTION

It is estimated that up to 50 million people sustain a traumatic brain injury (TBI) worldwide every year.[30] Of these, over 90 percent may be classified as “mild” based on the patient’s level of consciousness at presentation.[52] Such classification, however, is a misnomer as many of these patients are left with long-term adverse sequelae and disability. Despite the significant public health burden, our understanding of the trajectory of long-term recovery from mild TBI (mTBI) is limited. An improved understanding of the long-term sequelae after mTBI is important for ongoing clinical care of such patients, especially given the ongoing controversy over whether all mTBI patients recover within weeks versus a significant proportion having persisting ongoing disabling symptoms for months to years after the event.[4, 37, 38]

While traditionally defined as an initial Glasgow Coma Score of 13 to 15, loss of consciousness of less than 30 minutes, and post-traumatic amnesia no greater than 24 hours, the clinical spectrum of mTBI is broad, encompassing patients with subtle clinical signs and normal neuroimaging through those with clear evidence of injury on scans and/or more disabling neurological dysfunction.[2, 34, 46] Clinical outcome in mTBI can range from full and rapid recovery to chronic disabling symptoms.[30] A recent large cohort study found that 53% of patients with mTBI who had presented to US level 1 trauma centers reported continuing functional limitations at one year.[37, 38] These persisting symptoms are associated with mental health conditions such as post-traumatic stress disorder (PTSD), depression and suicide.[30] The impact of mTBI on quality of life (QoL) over time has been

relatively unstudied.[12]

Evidence for the breadth and duration of cognitive impairment following mTBI is mixed. Some studies report impairment spanning a wide range of domains; attention, working memory, episodic memory, information processing speed, general cognition,[21] whereas others emphasise deficits in attention and memory.[17] Early resolution of cognitive impairment (after one week[55] or one month[49]) is suggested by some studies whereas later resolution (after three[17, 23] or twelve[21] months) is suggested by others. Such differences in findings may reflect differing enrolment and severity characteristics.

The primary aim of this study was to examine longitudinal changes after mTBI, over an unusually long follow-up period of two years, in a comprehensive range of outcomes encompassing neurocognition, psychiatric symptoms, symptoms associated with post-concussion syndrome and quality of life.

MATERIALS AND METHODS

Participants

Seventy-three patients were recruited in the Emergency Department between 2012 and 2013 as part of the Acute Brain Injury Program at Addenbrooke's Hospital, Cambridge, United Kingdom which is a Major Trauma Center. Inclusion criteria included a Glasgow Coma Scale (GCS) score 13 to 15 on arrival to the Emergency Department (ED) and eligibility for Computed Tomography (CT) head due to head trauma according to NICE guidelines.[1] Patients with a past history of previous TBI, psychiatric or neurological disorder were

excluded. Participants were included in this analysis only if they attended a minimum of one follow up session (n = 61).

An additional 25 adults with extracranial injury only were recruited in the Emergency Department and had follow up assessments three months after injury and 99 healthy adults were also recruited. Control participants with suspected current or previous psychiatric disorder, neurological disorder, previous history of TBI or possible current TBI were excluded.

The Local Research Ethics Committee (NRES Committee Norfolk REC EE 0395) approved the study and written informed consent was obtained from all participants.

Neuropsychological assessment

Face-to-face assessment of health, quality of life (QoL) and cognitive function were completed at two weeks (median = 13 days, range = 6 to 23), three months (median = 92 days, range = 81 to 129), six months (median = 193 days, range = 169 to 253), one year (median = 375 days, range = 357 to 431), and two years (median = 736 days, range = 681 to 841) after injury. Very rarely (<2% of all instances) participants took home any questionnaires that were not completed during the face-to-face session due to time

constraints and returned them by post. The GOSE and all neurocognitive assessments were always performed face-to-face.

Health and QoL assessment

A battery of questionnaires was used to assess physical health, mental health and QoL: Alcohol Use Disorders Identification Test (AUDIT)[48] assessed drinking behaviour and alcohol-related problems; Beck Depression Inventory-II (BDI-II)[6] measured the magnitude of cognitive, behavioural and physiological symptoms of depression; PTSD Checklist-Civilian Version (PCL-C)[7] assessed PTSD symptoms; Brief Symptom Inventory-18 (BSI-18)[14] assessed psychological distress indicated by somatisation, depression and anxiety; Quality of Life after Brain Injury (QOLIBRI)[56] assessed TBI-specific QoL; Short Form-36 Health Survey (SF36)[8] assessed health-related QoL; Rivermead Post-Concussion Symptoms Questionnaire (RPQ)[24] assessed neurological/ somatic, emotional and cognitive post-concussion symptoms; Glasgow Outcome Scale Extended (GOSE)[58] structured interview categorised functional outcome (level of disability/recovery). All questionnaires have good reliability and validity and have been employed in previous TBI outcome studies.[39, 41, 52, 57]

Conventional dichotomisation of outcomes between GOSE 4 (Upper Severe Disability) and GOSE 5 (Lower Moderate Disability) was not appropriate for this cohort of patients, in whom an outcome would need to be back to or close to baseline to be classed as favourable. We therefore defined failure to achieve a Good Recovery (GOSE < 8) as an

unfavourable outcome. This approach is in keeping with current views on using sliding dichotomy or proportional odds methods in TBI studies.[26]

Cognitive functioning assessment

Cognitive functioning was assessed using classic phonemic (F A S words) and semantic (animal category words) verbal fluency tasks[19] and a selection of computerised tasks from the Cambridge Neuropsychological Test Automated Battery (CANTAB, cambridgecognition.com; administered on a Paceblade tablet). The battery of CANTAB tasks selected comprised: Motor Screening (MOT; screened for visual, motor and comprehension problems), Paired Associates Learning (PAL; episodic memory), Pattern Recognition Memory (PRM; visual recognition memory), Spatial Recognition Memory (SRM; spatial recognition memory), Spatial Span (SSP; visuospatial working memory capacity), Spatial Working Memory (SWM; retention and manipulation of visuospatial information, strategy use), Rapid Visual Information Processing (RVP; sustained attention), and Intra-Extra Dimensional Set

Shift (IED; visual discrimination, attentional flexibility). CANTAB tasks are sensitive to neurocognitive dysfunction following TBI.[29, 39-41]

Statistical analysis

Incidence of various health conditions at each time point was calculated using published cut-off scores; harmful drinking (AUDIT ≥ 8),[11] depression (BDI-II ≥ 14),[6] PTSD (PCL-C ≥ 33),[5] post-concussion syndrome (RPQ ≥ 3),[51] and GOSE < 8 . [26]

Our analyses used Latent Growth Curve (LGC) modelling techniques. LGC techniques are based on structural equation modelling and use repeated measures to estimate trajectories.[25, 50] Unlike traditional statistical techniques (e.g., repeated measures analysis of variance) they allow for the inclusion of participants with missing data (e.g., who dropped out or missed a testing session) enhancing the generalisability of findings. Inclusion of all data is particularly important in TBI research due to high attrition rates.[45] In addition to temporal changes at the group level, LGC modelling can also be used to explore individual differences in both temporal changes and initial status. Longitudinal changes in mean outcome scores were examined using LGC modelling in a structural equation modelling environment.[25, 50] The approach has been applied to modelling long-term changes in outcome after rehabilitation for TBI.[25] LGC modelling allowed for an exploration of the initial status of each measure (intercept), inter-individual differences in initial status (variance in intercept) and rate of change (variance in slope), and rate of group change over time (slope) using all available data. In cases where the linear LGC model was significant, further quadratic and cubic models were calculated; maximum likelihood estimation was used to determine which model provided the best fit for the data.

The distribution for CANTAB variables deviated from the normal distribution at some time points (as demonstrated by Shapiro-Wilk tests and P-P plots); prior to LGC analysis CANTAB scores were therefore transformed using square root transformation to help normalize the multivariate distributions (reverse transformation was first performed on variables for which lower scores equalled better performance). Growth curves were plotted with mTBI group means are plotted against trauma control norms (mean ± 1 SE) for health and QoL variables and against healthy control norms for the CANATB neurocognitive variables. Statistical analysis was performed using IBM SPSS Statistics (Version 25.0; Armonk, NY).

RESULTS

Table 1 presents participant demographic and injury characteristics. The majority of participants with mTBI (77%) were GCS 15 on arrival, although patients with GCS 13 (5%) and GCS 14 (18%) were also represented. Sixty-nine percent were male, and the age range 17 to 84 years (mean = 42.0 years, Standard Deviation (SD) ± 17.77). The estimated premorbid IQ range was 97 to 124 (mean = 113.72, SD ± 7.22) using the National Adult Reading Test (NART).[36] The trauma controls were 52% male with an age range 18 to 59 years (mean = 35.0 years, SD ± 12.42) and 99 healthy adults 49% male with an age range 18 to 70 years (mean = 36.3 years , SD ± 11.99).

Attendance over time

The number of participants in attendance at each follow-up session decreased over time from 56 (92% of the sample) at two weeks to 31 (51%) at two years (3 months, $n = 52$, 85%; 6 months, $n = 41$, 67%; 1 year, $n = 34$, 56%). Fifty two participants (85% of the sample) attended at least two follow-up sessions and 22 participants (36%) attended all five follow-up sessions (≥ 3 sessions, $n = 44$, 72%; ≥ 4 sessions, $n = 35$, 57%). No patients died during the study period to explain the loss to follow up. While exact causes of attrition were unable to be obtained the rate described is consistent with previous TBI studies.[44]

Incidence of health conditions and post-concussion symptoms

Figure 1 displays incidence rates of harmful drinking, depression, PTSD, post-concussion syndrome and disability and Figure 2 displays the percentage of participants reporting specific neurological, emotional and cognitive post-concussion symptoms at each time point (Fig 1 and 2). Incidence of harmful/hazardous drinking was as high at two years as it was at three months. Incidence of depression decreased steadily up to six months then rose to slightly higher than two-week levels. Incidences of PTSD, post-concussion syndrome and moderate-to-severe disability each decreased more steadily, but disability was noticeably lowest at six months.

Self-reported neurological symptoms tended to be most prevalent at two weeks; headaches, fatigue, dizziness, and sleep disturbance were reported most frequently. Incidence of headaches and dizziness decreased rapidly up to three months and then stayed relatively constant, although incidence of headaches was noticeably lowest at one year. Incidence of sleep disturbance decreased steadily up to six months then rose to two-week

levels. Incidence of fatigue decreased steadily to one year then rose again. Nausea, noise/light sensitivity and blurred/double vision did not fluctuate much over time.

Self-reported emotional symptoms were also most prevalent at two weeks with irritability and frustration being more common than depression and restlessness. Incidence of each emotional symptom decreased rapidly up to three months. After three months, incidence of depression remained relatively stable whereas incidences of restlessness, irritability and frustration rose again at six months.

Self-reported issues with cognition were reported by no less than one-fifth of participants at any time point. Incidences of concentration problems and slowed thinking declined from two weeks through two years whereas the opposite pattern was observed for memory problems which increased over time.

Longitudinal changes in outcome

The figures showing trajectories (Figure 3, Supplementary Figure 1 to 4) show a general trend for improvement over time, although the magnitude and rate of change varied substantially across variables. In the majority of cases, improvement appeared most rapid in the first three months after injury for health and QoL variables and in the first six months after injury for cognitive variables.

The trajectory of key mean scores to two years post injury can be seen in Figure 3. In the patient cohort, scores on GOSE, BSI-19, and BDI were worse, and harmful drinking significantly more common than control norms at least until three months, and remained so

for some outcomes at every follow up time point. Any change in these variables were not significant as demonstrated by non-significant linear LGC model slopes (Table 2). The rate of improvement in QOLIBRI was rapid for the first three months and then decelerated with QOLIBRI not reaching trauma control norms until two years after injury (Fig 3). The best fit for this trajectory was quadratic with a significant model slope ($p = 0.034$). The Rivermead post-concussion symptoms appeared to improve rapidly up to three months and then plateaued up to one year. The cubic LGC model provided the best fit for improvements in post-concussion symptoms demonstrated by a significant cubic LGC slope and significant differences between the -2LL for the quadratic and cubic models. Detailed results for SF-36, and the CANTAB neurocognitive battery may be found in the Supplementary Material (Supplementary Figure 1 to 4, Table 2).

DISCUSSION

This study examined recovery from mTBI over two years, and found that ongoing symptoms were common for the entire period. Importantly, it is the first study to report functional, quality of life and neurocognitive outcomes over such a long time period. Although improvement was rapid early on (before plateauing) for post-concussion symptoms (significant improvement up to three months) and GOSE, scores fell below trauma control norms at every time point. The overall persistence of symptoms was high; two years after injury, one-third of the sample reported headaches, fatigue or sleep disturbances, and one-fifth had limitations with a GOSE <8. Incidence of unfavourable outcome at earlier time points was similar to figures reported elsewhere.[12] Consistent with prior research, symptoms commonly associated with post-concussion syndrome including headaches, fatigue, dizziness and sleep disturbance were the most common symptoms.[43] Nausea, noise/light sensitivity, and blurred/double vision were less common but did not diminish over time.

Mental health related symptoms were common; incidences of depression (16%), PTSD (9%) and harmful drinking (35%) were above national averages even two years post-injury.[15] Feelings of irritability and frustration fell from 40% at two weeks to 15% at two years. Incidence of PTSD following mTBI was similar to that reported elsewhere whereas incidence of depression was higher here.[20] Depression and sleep disturbance followed a similar pattern of steady decrease in incidence up to six months before rising to two week levels, suggesting that the “neurological” symptoms of the RPQ may be a consequence of depression.[22] Depression is one of the most common psychiatric sequelae in survivors of TBI, and its presence is associated with impaired functional and cognitive recovery, and

increased disability.[3] The exact mechanisms are unclear but are thought to be (at least in part) secondary to altered functional connectivity of networks associated with emotional regulation.[35]

Depression, psychological distress and harmful drinking scores were poorer than trauma control norms at all time points (although longitudinal changes were not significant). Interestingly, the two measures of depression revealed different two-week incidence rates; reporting depressed mood on the RPQ (25%) was more common than being classified as depressed on the BDI-II (14%). The act of reporting other symptoms during completion of the RPQ may have enhanced perception of current low mood highlighting the importance of considering contextual factors when analysing questionnaire data.[54]

Ongoing lowered QoL was predominately due to emotional rather than physical health factors. Physical-related, social-related and TBI-specific QoL improved rapidly early-on and tended to reach trauma control norms by three months. Bodily pain scores were better than trauma control norms, which is unsurprising given the nature of their extracranial injuries. Emotional-related QoL improved steadily through all time points, and like TBI-specific QoL (which combined emotional, physical, social and cognitive factors), did not reach trauma control norms until one or two year(s). Temporal changes in QoL after mTBI have been little studied, although links between emotional health (depression and PTSD) following mTBI and lowered QoL have been reported.[20]

Cognitive problems—slowed thinking, poor concentration or memory problems—were reported by no less than one fifth of participants at any given time point. Self-reports

were corroborated by actual performance on the cognitive tasks in almost every case. Concentration problems were reported by approximately one-third of the sample up to one year before dropping considerably; this was mirrored by performance on the sustained attention task on which improvement (RVP A prime) was slow up to one year then accelerated. Self-reports of slowed thinking decreased over time (from 31% to 26%) with a slight increase between six months and one year; this was reflected in latency scores which improved steadily over time (RVP latency) or improved rapidly up to six months before worsening (PRM latency). Memory problems were reported more often as time progressed and by more than approximately one third of the sample at all time points; similarly, performance on the memory tasks tended to remain lower than healthy control norms. Consistent with other literature, memory issues were more common than poor concentration and slowed thinking,[28] and reductions in response speed[32] and attention deficits[31] following mTBI are particularly common in the first few months after injury but can continue to one year or later.[42]

Temporal changes were also observed for the majority of cognitive domains (all except working memory). Improvement in semantic verbal fluency was steady over time, whereas improvement in phonemic verbal fluency was rapid up to six months then slower. Phonemic verbal fluency was below healthy control norms only at two weeks. Greater semantic relative to phonemic verbal fluency deficits in mild-to-moderate TBI have been reported elsewhere.[18] Attentional flexibility and working memory accuracy scores were poorer than healthy control norms at all time points. Attentional flexibility (IED pre-ED errors) improvement was rapid until three months then decelerated but working memory improvement was non-significant over time. Counter to the abrupt nature of strategy

discovery reported elsewhere,[27] strategy use (SWM strategy) improved linearly over time but never reached healthy control norms. Evidence for working memory deficits following mTBI is mixed; working memory deficits have been reported in some studies,[13, 55] but other studies have found no performance differences between mTBI patients and controls despite differential brain activation patterns with increased working memory load.[33]

Future studies using imaging techniques such as diffusion tensor imaging (DTI), functional magnetic resonance imaging, and spectroscopy may provide valuable insight into the causes of prolonged issues in mTBI. For example, DTI characterises the diffusion of water molecules which is influenced by the microstructural organization of tissues, particularly in white matter. Measuring longitudinal changes from DTI may offer insights into structural connectivity changes and their relationship to outcome. Interestingly, the cognitive measures that worsened over time in our sample (memory and processing speed) are also those linked with decreases in white matter integrity over time in moderate-to-severe TBI.[40] Recent studies have linked diffuse axonal injury in the limbic system with PTSD in mTBI.[47] Indeed, many of the cognitive deficits and mental health symptoms experienced following mTBI may be caused by damaged attentional control pathways.[16] For therapies to be targeted effectively and with the greatest gains in outcome, further studies should establish the causal structure of prolonged problems. Targeted sleep therapy, for example, may not only provide direct resolution of sleep disturbances but may also provide indirect relief of co-occurring problems such as those with cognition and emotion. Similarly, targeted therapy for depression may also improve cognitive

functioning.[10] In addition, outcome may be influenced by the presence of extra-cranial injury, and larger studies may allow the effects of this to be fully elucidated.[9]

In this study attendance rate fell from 92% to 51% between two weeks and two years after injury and only 36% of our sample of adults with mTBI attended all five testing occasions. By employing LGC modelling instead of more traditional analysis techniques we avoided substantial loss of power, bias and lowered generalisability because data from participants that missed testing sessions could be included.

Limitations to this study included the relatively small sample size and lack of control for practice effects and some premorbid factors. A significant proportion of the variance in all LGC models was left unexplained and the sample size prevented an investigation of predictive variables via the addition of various covariates. Despite efforts to exclude participants with diagnosed psychiatric conditions, premorbid levels of depression, anxiety and substance abuse, which are known predictors of post-TBI functioning,[42] were not recorded. It is therefore impossible to separate the apparent consequences of mTBI observed here from existing characteristics. Practice effects are an unavoidable consequence of repeated cognitive testing, especially on novel executive tasks[53] and tasks promoting strategy formation,[5] and are observed on many of the CANTAB tasks employed here. Unfortunately, normative data, separated by age and ability level, on serial CANTAB testing is not available. However, our inclusion of control norms and self-report measures of cognition help to overcome some of the difficulty with distinguishing genuine improvement from task familiarity. At least some influence of recovery over practice was suggested by performance rarely reaching normal levels, even after multiple testing sessions and

considerable recovery time, and by corroboration of performance improvements by participants' self-reports.

The elevated level and duration of cognitive and emotional issues observed here may be attributable to our use of techniques which do not necessitate the removal of cases with missing data. If the nature of the missing data is not random, such that non-attenders tend to have extreme outcomes limiting their ability or motivation to attend, ongoing difficulties will be underestimated.[45]

CONCLUSION

This study demonstrates that, for some individuals, recovery from mTBI occurs over a long timescale, and is much more protracted than usually believed. In particular, problems with emotional well-being, alcohol use and cognition, as well as symptoms such as headaches and sleep disturbance, are common after mTBI and often long-standing. Further LGC modelling in larger samples and over longer time frames will enable exploration of casual factors whilst providing the power to detect potential individual differences in the rate of change over time. The findings point to the need for long-term management of mTBI targeting subgroups with persisting decreases in cognition, mental health problems, and functional limitations.

Declarations

Acknowledgements: The NIHR/Wellcome Trust Cambridge Clinical Research Facility provided nursing support to aid in the patient recruitment.

Funding: This work was partially funded by the European Commission under the 7th Framework Programme (FP7-270259-TBIcare), the UK National Institute of Health Research (NIHR) Biomedical Research Centre at Cambridge and the Technology Platform funding provided by the UK Department of Health. PJH was supported by the NIHR (Research Professorship and Cambridge BRC). JPP received funding from the Academy of Finland (grant #17379), Government's Special Financial Transfer tied to academic research in Health Sciences (Finland), the Emil Aaltonen Foundation, the Finnish Brain Foundation and the Maire Taponen Foundation. BJS receives funding from the NIHR Brain Injury MedTech Co-operative, Cambridge and the NIHR Cambridge Biomedical Research Centre (Mental Health Theme). DKM was supported by the NIHR through the NIHR Cambridge Biomedical Research Centre grant and a Senior Investigator Award. VFJN was funded by a Health Foundation/Academy of Medical Sciences Clinician Scientist Fellowship.

Conflicts of interest: BJS consults for Cambridge Cognition and Greenfield BioVentures and Cassava Sciences. VFJN holds a grant from Roche Pharmaceuticals for a project not related to this study. No other authors have any conflicts of interests to declare.

Author Contributions: ELC, DKM and VFJN conceptualized and designed the study. ELC, JGO, FF, AEM, PJAH, OT, JPP, LW, BJS, DKM and VFJN were involved with data acquisition and analysis. ELC and VFJN drafted the manuscript. All authors edited the manuscript.

Data availability statement: Anonymised data used for the analyses in this study are available from the corresponding author upon reasonable request (ORCID ID <https://orcid.org/0000-0001-6044-9035>).

REFERENCES

1. (2014) Head injury: assessment and early management NICE Clinical Guideline [CG176]:<https://www.nice.org.uk/guidance/cg176>
2. ACRM, Committee MTBI (1993) Definition of mild traumatic brain injury. *J Head Trauma Rehabil* 8:86-87
3. Albrecht JS, Lydecker A, Peters M, Rao V (2020) Treatment of Depression Following Traumatic Brain Injury Reduces Risk of Neuropsychiatric Outcomes. *J Neurotrauma*
4. Andrikopoulos J (2019) The Term Mild in Mild Traumatic Surgery and the Representation of Outcomes. *JAMA Neurol*
5. Basso MR, Bornstein RA, Lang JM (1999) Practice effects on commonly used measures of executive function across twelve months. *Clin Neuropsychol* 13:283-292
6. Beck AT, Steer RA, Ball R, Ranieri W (1996) Comparison of Beck Depression Inventories -IA and -II in psychiatric outpatients. *J Pers Assess* 67:588-597
7. Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA (1996) Psychometric properties of the PTSD Checklist (PCL). *Behav Res Ther* 34:669-673
8. Brazier JE, Harper R, Jones NM, O'Cathain A, Thomas KJ, Usherwood T, Westlake L (1992) Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ* 305:160-164
9. Carroll EL, Manktelow AE, Outtrim JG, Chatfield D, Forsyth F, Hutchinson PJA, Tenovuo O, Posti JP, Wilson L, Sahakian BJ, Menon DK, Newcombe VFJ (2020) Influence of Concomitant Extracranial Injury on Functional and Cognitive Recovery From Mild Versus Moderate to Severe Traumatic Brain Injury. *J Head Trauma Rehabil*
10. Chamelian L, Feinstein A (2006) The effect of major depression on subjective and objective cognitive deficits in mild to moderate traumatic brain injury. *J Neuropsychiatry Clin Neurosci* 18:33-38
11. Conigrave KM, Hall WD, Saunders JB (1995) The AUDIT questionnaire: choosing a cut-off score. *Alcohol Use Disorder Identification Test. Addiction* 90:1349-1356
12. de Koning ME, Scheenen ME, van der Horn HJ, Timmerman ME, Hageman G, Roks G, Spikman JM, van der Naalt J (2017) Prediction of work resumption and sustainability up to 1 year after mild traumatic brain injury. *Neurology* 89:1908-1914
13. Dean PJ, Sterr A (2013) Long-term effects of mild traumatic brain injury on cognitive performance. *Front Hum Neurosci* 7:30
14. Derogatis LR, Melisaratos N (1983) The Brief Symptom Inventory: an introductory report. *Psychol Med* 13:595-605
15. Drummond C, McBride O, Fear N, Fuller E (2016) Mental health and wellbeing in England: Adult Psychiatric Morbidity Survey 2014. NHS Digital
16. Etkin A, Buchel C, Gross JJ (2015) The neural bases of emotion regulation. *Nat Rev Neurosci* 16:693-700
17. Flynn FG (2010) Memory impairment after mild traumatic brain injury. *Continuum (Minneapolis)* 16:79-109
18. Goldstein FC, Levin HS, Roberts VJ, Goldman WP, Kalechstein AS, Winslow M, Goldstein SJ (1996) Neuropsychological effects of closed head injury in older adults: A comparison with Alzheimer's disease. *Neuropsychology* 10:147-154
19. Goodglass H, Kaplan E (1983) Boston diagnostic aphasia examination booklet. Lea & Febiger;
20. Haagsma JA, Scholten AC, Andriessen TM, Vos PE, Van Beeck EF, Polinder S (2015) Impact of depression and post-traumatic stress disorder on functional outcome and health-related quality of life of patients with mild traumatic brain injury. *J Neurotrauma* 32:853-862
21. Heitger MH, Jones RD, Dalrymple-Alford JC, Frampton CM, Ardagh MW, Anderson TJ (2006) Motor deficits and recovery during the first year following mild closed head injury. *Brain Inj* 20:807-824

22. Iverson GL (2006) Misdiagnosis of the persistent postconcussion syndrome in patients with depression. *Arch Clin Neuropsychol* 21:303-310
23. Karr JE, Areshenkoff CN, Garcia-Barrera MA (2014) The neuropsychological outcomes of concussion: a systematic review of meta-analyses on the cognitive sequelae of mild traumatic brain injury. *Neuropsychology* 28:321-336
24. King NS, Crawford S, Wenden FJ, Moss NE, Wade DT (1995) The Rivermead Post Concussion Symptoms Questionnaire: a measure of symptoms commonly experienced after head injury and its reliability. *J Neurol* 242:587-592
25. Kozlowski AJ, Pretz CR, Dams-O'Connor K, Kreider S, Whiteneck G (2013) An introduction to applying individual growth curve models to evaluate change in rehabilitation: a National Institute on Disability and Rehabilitation Research Traumatic Brain Injury Model Systems report. *Arch Phys Med Rehabil* 94:589-596
26. Lingsma HF, Roozenbeek B, Steyerberg EW, Murray GD, Maas AI (2010) Early prognosis in traumatic brain injury: from prophecies to predictions. *Lancet Neurol* 9:543-554
27. Lowe C, Rabbitt P (1998) Test/re-test reliability of the CANTAB and ISPOCD neuropsychological batteries: theoretical and practical issues. Cambridge Neuropsychological Test Automated Battery. International Study of Post-Operative Cognitive Dysfunction. *Neuropsychologia* 36:915-923
28. Lundin A, de Boussard C, Edman G, Borg J (2006) Symptoms and disability until 3 months after mild TBI. *Brain Inj* 20:799-806
29. Lunter CM, Carroll EL, Housden C, Outtrim J, Forsyth F, Rivera A, Maimaris C, Boyle A, Sahakian BJ, Menon DK, Newcombe VF (2019) Neurocognitive testing in the emergency department: A potential assessment tool for mild traumatic brain injury. *Emerg Med Australas* 31:355-361
30. Maas AIR, Menon DK, Adelson PD, Andelic N, Bell MJ, Belli A, Bragge P, Brazinova A, Buki A, Chesnut RM, Citerio G, Coburn M, Cooper DJ, Crowder AT, Czeiter E, Czosnyka M, Diaz-Arrastia R, Dreier JP, Duhaime AC, Ercole A, van Essen TA, Feigin VL, Gao G, Giacino J, Gonzalez-Lara LE, Gruen RL, Gupta D, Hartings JA, Hill S, Jiang JY, Ketharanathan N, Kompanje EJO, Lanyon L, Laureys S, Lecky F, Levin H, Lingsma HF, Maegele M, Majdan M, Manley G, Marsteller J, Mascia L, McFadyen C, Mondello S, Newcombe V, Palotie A, Parizel PM, Peul W, Piercy J, Polinder S, Puybasset L, Rasmussen TE, Rossaint R, Smielewski P, Soderberg J, Stanworth SJ, Stein MB, von Steinbuechel N, Stewart W, Steyerberg EW, Stocchetti N, Synnot A, Te Ao B, Tenovuo O, Theadom A, Tibboel D, Videtta W, Wang KKW, Williams WH, Wilson L, Yaffe K, In TP, Investigators (2017) Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol* 16:987-1048
31. Maruta J, Suh M, Niogi SN, Mukherjee P, Ghajar J (2010) Visual tracking synchronization as a metric for concussion screening. *J Head Trauma Rehabil* 25:293-305
32. Mathias JL, Beall JA, Bigler ED (2004) Neuropsychological and information processing deficits following mild traumatic brain injury. *J Int Neuropsychol Soc* 10:286-297
33. McAllister TW, Sparling MB, Flashman LA, Guerin SJ, Mamourian AC, Saykin AJ (2001) Differential working memory load effects after mild traumatic brain injury. *Neuroimage* 14:1004-1012
34. Menon DK, Schwab K, Wright DW, Maas AI, Demographics, Clinical Assessment Working Group of the I, Interagency Initiative toward Common Data Elements for Research on Traumatic Brain I, Psychological H (2010) Position statement: definition of traumatic brain injury. *Arch Phys Med Rehabil* 91:1637-1640
35. Moreno-Lopez L, Sahakian BJ, Manktelow A, Menon DK, Stamatakis EA (2016) Depression following traumatic brain injury: A functional connectivity perspective. *Brain Inj* 30:1319-1328
36. Nelson HE (1982) National Adult Reading Test. NFER-Nelson

37. Nelson LD, Levin HS, McCrea MA (2019) The Term Mild in Mild Traumatic Surgery and the Representation of Outcomes-Reply. *JAMA Neurol*
38. Nelson LD, Temkin NR, Dikmen S, Barber J, Giacino JT, Yuh E, Levin HS, McCrea MA, Stein MB, Mukherjee P, Okonkwo DO, Diaz-Arrastia R, Manley GT, and the T-TBII, Adeoye O, Badjatia N, Boase K, Bodien Y, Bullock MR, Chesnut R, Corrigan JD, Crawford K, Mis, Duhaime AC, Ellenbogen R, Feeser VR, Ferguson A, Foreman B, Gardner R, Gaudette E, Gonzalez L, Gopinath S, Gullapalli R, Hemphill JC, Hotz G, Jain S, Korley F, Kramer J, Kreitzer N, Lindsell C, Machamer J, Madden C, Martin A, McAllister T, Merchant R, Noel F, Palacios E, Perl D, Puccio A, Rabinowitz M, Robertson CS, Rosand J, Sander A, Satris G, Schnyer D, Seabury S, Sherer M, Taylor S, Toga A, Valadka A, Vassar MJ, Vespa P, Wang K, Yue JK, Zafonte R (2019) Recovery After Mild Traumatic Brain Injury in Patients Presenting to US Level I Trauma Centers: A Transforming Research and Clinical Knowledge in Traumatic Brain Injury (TRACK-TBI) Study. *JAMA Neurol*
39. Newcombe V, Chatfield D, Outtrim J, Vowler S, Manktelow A, Cross J, Scoffings D, Coleman M, Hutchinson P, Coles J, Carpenter TA, Pickard J, Williams G, Menon D (2011) Mapping traumatic axonal injury using diffusion tensor imaging: correlations with functional outcome. *PLoS One* 6:e19214
40. Newcombe VF, Correia MM, Ledig C, Abate MG, Outtrim JG, Chatfield D, Geeraerts T, Manktelow AE, Garyfallidis E, Pickard JD, Sahakian BJ, Hutchinson PJ, Rueckert D, Coles JP, Williams GB, Menon DK (2016) Dynamic Changes in White Matter Abnormalities Correlate With Late Improvement and Deterioration Following TBI: A Diffusion Tensor Imaging Study. *Neurorehabil Neural Repair* 30:49-62
41. Newcombe VF, Outtrim JG, Chatfield DA, Manktelow A, Hutchinson PJ, Coles JP, Williams GB, Sahakian BJ, Menon DK (2011) Parcellating the neuroanatomical basis of impaired decision-making in traumatic brain injury. *Brain* 134:759-768
42. Ponsford J, Tweedly L, Lee N, Taffe J (2012) Who responds better? Factors influencing a positive response to brief alcohol interventions for individuals with traumatic brain injury. *J Head Trauma Rehabil* 27:342-348
43. Ponsford JL, Downing MG, Olver J, Ponsford M, Acher R, Carty M, Spitz G (2014) Longitudinal follow-up of patients with traumatic brain injury: outcome at two, five, and ten years post-injury. *J Neurotrauma* 31:64-77
44. Richter S, Stevenson S, Newman T, Wilson L, Maas AIR, Nieboer D, Lingsma H, Steyerberg EW, Newcombe VFJ (2020) Study Design Features Associated with Patient Attrition in Studies of Traumatic Brain Injury: A Systematic Review. *J Neurotrauma*
45. Richter S, Stevenson S, Newman T, Wilson L, Menon DK, Maas AIR, Nieboer D, Lingsma H, Steyerberg EW, Newcombe VFJ (2019) Handling of Missing Outcome Data in Traumatic Brain Injury Research: A Systematic Review. *J Neurotrauma*
46. Ruff RM, Iverson GL, Barth JT, Bush SS, Broshek DK, Policy NAN, Planning C (2009) Recommendations for diagnosing a mild traumatic brain injury: a National Academy of Neuropsychology education paper. *Arch Clin Neuropsychol* 24:3-10
47. Santhanam P, Teslovich T, Wilson SH, Yeh PH, Oakes TR, Weaver LK (2019) Decreases in White Matter Integrity of Ventro-Limbic Pathway Linked to Post-Traumatic Stress Disorder in Mild Traumatic Brain Injury. *J Neurotrauma* 36:1093-1098
48. Saunders JB, Aasland OG, Babor TF, de la Fuente JR, Grant M (1993) Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption--II. *Addiction* 88:791-804
49. Schretlen DJ, Shapiro AM (2003) A quantitative review of the effects of traumatic brain injury on cognitive functioning. *Int Rev Psychiatry* 15:341-349
50. Shek DTL, Ma CMS (2011) Longitudinal Data Analyses Using Linear Mixed Models in SPSS: Concepts, Procedures and Illustrations. *Thescientificworldjo* 11:42-76

51. Sigurdardottir S, Andelic N, Roe C, Jerstad T, Schanke AK (2009) Post-concussion symptoms after traumatic brain injury at 3 and 12 months post-injury: a prospective study. *Brain Inj* 23:489-497
52. Steyerberg EW, Wiegers E, Sewalt C, Buki A, Citerio G, De Keyser V, Ercole A, Kunzmann K, Lanyon L, Lecky F, Lingsma H, Manley G, Nelson D, Peul W, Stocchetti N, von Steinbuechel N, Vande Vyvere T, Verheyden J, Wilson L, Maas AIR, Menon DK, Participants C-T, Investigators (2019) Case-mix, care pathways, and outcomes in patients with traumatic brain injury in CENTER-TBI: a European prospective, multicentre, longitudinal, cohort study. *Lancet Neurol* 18:923-934
53. Stuss DT (2011) Traumatic brain injury: relation to executive dysfunction and the frontal lobes. *Curr Opin Neurol* 24:584-589
54. Tourangeau R (2004) Survey research and societal change. *Annu Rev Psychol* 55:775-801
55. Vanderploeg RD, Curtiss G, Belanger HG (2005) Long-term neuropsychological outcomes following mild traumatic brain injury. *J Int Neuropsychol Soc* 11:228-236
56. von Steinbuechel N, Wilson L, Gibbons H, Hawthorne G, Hofer S, Schmidt S, Bullinger M, Maas A, Neugebauer E, Powell J, von Wild K, Zitnay G, Bakx W, Christensen AL, Koskinen S, Sarajuuri J, Formisano R, Sasse N, Truelle JL, Force QT (2010) Quality of Life after Brain Injury (QOLIBRI): scale development and metric properties. *J Neurotrauma* 27:1167-1185
57. Voormolen DC, Haagsma JA, Polinder S, Maas AIR, Steyerberg EW, Vulekovic P, Sewalt CA, Gravesteijn BY, Covic A, Andelic N, Plass AM, von Steinbuechel N (2019) Post-Concussion Symptoms in Complicated vs. Uncomplicated Mild Traumatic Brain Injury Patients at Three and Six Months Post-Injury: Results from the CENTER-TBI Study. *J Clin Med* 8
58. Wilson JT, Pettigrew LE, Teasdale GM (1998) Structured interviews for the Glasgow Outcome Scale and the extended Glasgow Outcome Scale: guidelines for their use. *J Neurotrauma* 15:573-585

FIGURE LEGENDS

Figure 1. Incidence of health conditions for patients with mild traumatic brain injury —harmful drinking (A; AUDIT ≥ 8), depression (A; BDI-II ≥ 14), PTSD (A; PCL-C ≥ 33), PCS (B; RPQ ≥ 3), disability (B; GOSE < 8)—at each time point. Abbreviations: PCS = post-concussion syndrome; PTSD = post-traumatic stress disorder; AUDIT = Alcohol Use Disorders Identification Test; BDI = Beck Depression Inventory; PCL-C = PTSD Checklist-Civilian Version; RPQ = Rivermead Post-Concussion Symptoms Questionnaire; GOSE = Glasgow Outcome Score Extended.

Figure 2. Incidence of self-reported post-concussion symptoms for patients with mild traumatic brain injury—neurological (A), emotional (B), cognitive (B)—at each time point.

Figure 3. Mean health scores for patients with mild traumatic brain injury: Glasgow Outcome Score Extended (GOSE) (A), Rivermead Post-Concussion Questionnaire number of symptoms (RPQ) (B), Quality of life after brain injury (QOLIBRI) (C), Psychological distress (Brief Symptom Inventory-18, BSI-18) (D), depression (Beck depression inventory-II (BDI-II)) and (E) harmful drinking (Alcohol Use Disorders Identification Test (AUDIT)) (F). Error bars represent standard error. Dotted lines represent mean (± 1 SE) scores in trauma controls at three months after extracranial injury.

Table 1: Participant characteristics across the groups of healthy volunteers, trauma controls and patients with mild traumatic brain injury.

Participant Characteristic	Healthy volunteer controls (n = 99)	Trauma controls (n = 25)	mTBI (n = 61)	Kruskal-Wallis ^a or Chi-Square ^b		
				df	H or χ^2	p
Age, mean (SD), years	36 (12)	35 (12)	42(18)	2	3.2 ^a	0.202
Sex, n (%) female	50 (51)	12 (48)	19(31)	2	5.77 ^b	0.056
GCS, n(%)						
15	-	-	47(77)			
14			11(18)			
13			3(5)			
AIS Total, median (range)	-	1 (1-5)	4 (1-16)			
AIS Extracranial, median (range)	-	1 (1-5)	2(1-12)			
Marshall Score, n(%)						
I	-	-	53(87)			
II	-	-	7(11.4)			
III	-	-	0			
IV	-	-	0			
V	-	-	0			
VI	-	-	1(1.6)			
Mechanism of Injury						
Road Traffic Collision	-	7(28)	14(23)	6	15.44 ^b	0.017
Fall	-	9(36)	21(34)			
Violence or assault	-	0	6(10)			
Blow to head (not assault)	-	0	7(12)			
Rugby injury	-	2(8)	4(7)			
Cycling or Horse riding	-	3(12)	9(15)			
Blow to body (not head)	-	4(16)	0			
Education, n(%)						
Low	41 (43)	10 (40)	19(31)	4	2.41 ^b	0.660
Middle	26 (27)	5 (20)	17(28)			
High	29 (30)	10 (40)	25(41)			

Abbreviations: mTBI, mild traumatic brain injury; Glasgow Coma Scale; AIS, Abbreviated Injury Scale; SD, standard deviation; n, number.

Table 2. Latent Growth Curve modelling results for patients with mild traumatic brain injury

Outcome measure	LGC model	group intercept			group slope			variance of intercept		variance of slope		covariance of intercept &		residual	
		df	F	P	df	F	p	Z	p	Z	p	Z	p	Z	p
AUDIT	linear	1, 58.78	101.37	< 0.001	1, 24.78	1.60	0.217	5.00	< 0.001	0.80	0.427	-1.44	0.151	7.14	< 0.001
DBI-II	linear	1, 57.62	67.29	< 0.001	1, 36.46	0.92	0.346	3.81	< 0.001	0.39	0.698	-0.55	0.585	7.62	< 0.001
BSI-18	linear	1, 50.69	56.35	< 0.001	1, 24.05	3.23	0.085	3.19	0.001	0.2	0.771	-1.15	0.250	7.18	< 0.001
RPQ	cubic	1, 75.71	85.69	< 0.001	1, 123.26	4.93	0.028	4.60	< 0.001	1.35	0.176	-1.99	0.047	7.62	< 0.001
GOSE outcome	linear	1, 56.54	3576.	< 0.001	1, 37.23	0.90	0.350	2.12	0.024	0.6	0.507	-0.41	0.681	6.19	< 0.001
QOLIBRI	quadrati	1, 68.84	1143.7	< 0.001	1, 130.27	4.58	0.034	4.48	< 0.001	1.55	0.121	-1.98	0.048	7.30	< 0.001
Physical	linear	1, 59.54	997.20	< 0.001	1, 36.05	2.68	0.002	4.28	< 0.001	1.61	0.109	-2.33	0.020	7.24	< 0.001
Social	cubic	1,	469.02	< 0.001	1, 123.71	10.65	0.001	3.24	< 0.001	1.08	0.283	-1.89	0.058	7.53	< 0.001
Role physical	cubic	1, 86.14	74.94	< 0.001	1, 111.94	6.98	0.009	4.01	< 0.001	1.25	0.212	-1.90	0.058	7.18	< 0.001
Role emotional	linear	1, 50.57	531.99	< 0.001	1, 31.18	3.99	0.055	3.19	0.001	0.46	0.645	-1.45	0.146	7.20	< 0.001
Emotional well-	linear	1, 24.92	1518.3	< 0.001	1, 3.43	10.62	0.039	2.17	0.030	-	-	-0.23	0.821	3.44	0.001
Energy & vitality	linear	1, 45.38	631.71	< 0.001	1, 101.25	10.48	0.002	2.97	0.003	-	-	-0.15	0.883	7.58	< 0.001
Bodily pain	cubic	1, 92.97	243.69	< 0.001	1, 108.79	16.84	<	3.98	< 0.001	1.37	0.169	-1.43	0.152	7.03	< 0.001
General health	linear	1, 56.94	803.64	< 0.001	1, 31.18	0.06	0.805	4.52	< 0.001	0.20	0.840	1.09	0.277	7.36	< 0.001
Phonemic VF	quadrati	1, 65.24	896.60	< 0.001	1, 129.42	10.92	0.001	4.33	< 0.001	0.32	0.751	0.42	0.672	7.10	< 0.001
Semantic VF	linear	1, 53.17	1386.6	< 0.001	1, 143.03	4.90	0.028	3.25	0.001	-	-	0.89	0.375	8.06	< 0.001
PAL errors	linear	1, 68.17	9356.7	< 0.001	1, 156.98	0.28	0.601	3.64	< 0.001	-	-	-	-	8.48	< 0.001
PAL stages	linear	1, 37.76	5239.1	< 0.001	1, 147.42	0.02	0.900	2.60	0.009	-	-	0.43	0.670	8.15	< 0.001
SRM latency	linear	1, 57.33	4351.5	< 0.001	1, 147.95	1.52	0.219	3.20	0.001	-	-	-	-	8.15	< 0.001
SRM percent	quadrati	1, 86.80	14508.	< 0.001	1, 164.66	6.64	0.011	3.46	0.001	-	-	-0.19	0.854	8.85	< 0.001
PRM latency	cubic	1, 93.54	4385.6	< 0.001	1, 111.46	6.46	0.012	3.81	< 0.001	1.71	0.088	-2.90	0.004	7.19	< 0.001
PRM percent	linear	1, 54.43	19204.	< 0.001	1, 143.52	2.00	0.160	3.88	< 0.001	-	-	-0.96	0.338	8.43	< 0.001
SSP span length	linear	1, 59.13	4174.2	< 0.001	1, 36.12	0.27	0.607	4.54	< 0.001	1.45	0.147	0.26	0.797	7.33	< 0.001
SWM between	linear	1, 60.58	1663.9	< 0.001	1, 0.38	34.52	0.540	4.77	< 0.001	1.65	0.098	-0.49	0.627	7.37	< 0.001
SWM within	linear	1, 57.38	4865.5	< 0.001	1, 25.85	0.13	0.722	4.08	< 0.001	1.95	0.051	-2.87	0.004	7.33	< 0.001
SWM strategy	linear	1, 59.44	787.49	< 0.001	1, 36.21	7.88	0.008	4.77	< 0.001	1.19	0.234	-0.43	0.664	7.44	< 0.001
RVP A prime	quadrati	1, 67.20	67125.	< 0.001	1, 139.08	6.55	0.012	4.61	< 0.001	-	-	0.92	0.360	8.14	< 0.001
RVP latency	linear	1, 47.46	2410.0	< 0.001	1, 93.46	4.99	0.028	3.36	0.001	-	-	-	-	6.65	< 0.001
IED pre-ED errors	quadrati	1, 68.22	999.26	< 0.001	1, 73.72	12.23	0.001	4.09	< 0.001	-	-	-	-	6.28	< 0.001
IED ED errors	linear	1, 0.01	2890.5	0.944	1, 0.03	2.90	0.917	0.04	0.970	-	-	-	-	0.24	< 0.001
IED reversal	linear	1,	6861.4	< 0.001	1, 147.47	0.13	0.719	-	-	-	-	-	-	8.13	< 0.001

Abbreviations: LGC = latent growth curve; PCS = post-concussion syndrome; GOSE = Glasgow Outcome Score Extended; QoL = quality of life; VF = verbal fluency; PAL = paired associates learning; SRM = spatial recognition memory; PRM = pattern recognition memory; SSP = spatial span; SWM = spatial working memory; RVP = rapid visual information processing; IED = inter-extra dimensional set shifting; ED = extra dimensional.